

# Parasites of four ornamental fish from the Chumucuí River (Bragança, Pará, Brazil)

Parasitas de quatro peixes ornamentais do Rio Chumucuí (Bragança-Pará, Brasil)

Rodrigo Yudi Fujimoto<sup>1\*</sup>; Zaira Monik Nunes de Barros<sup>2</sup>; Adjalbas Nunes Marinho-Filho<sup>2</sup>;  
Daniel Guerreiro Diniz<sup>3</sup>; Jorge Costa Eiras<sup>4</sup>

<sup>1</sup>Embrapa Tabuleiros Costeiros, Aracajú, SE, Brasil

<sup>2</sup>Laboratório de Ictioparasitologia e Piscicultura, Universidade Federal do Pará – UFPA, Bragança, PA, Brasil

<sup>3</sup>Laboratório de Investigações em Neurodegeneração e Infecção, Instituto de Ciências Biológicas, Hospital Universitário João de Barros Barreto, Universidade Federal do Pará – UFPA, Belém, PA, Brasil

<sup>4</sup>Laboratório Associado – CIMAR, Departamento de Biologia, Centro Interdisciplinar de Investigação Marinha e Ambiental – CIIMAR, Faculdade de Ciências, Universidade do Porto – U. Porto, Porto, Portugal

Received March 12, 2012

Accepted October 29, 2012

## Abstract

The objective of the present study was to evaluate the parasite fauna of four species of ornamental fish collected in the Chumucuí River, municipality of Bragança, Pará, Brazil. From June 2006 to December 2007. Fishes (n=307) belonging to four species were collected, including 23 specimens of *Moenkhausia sanctaefilomenae* (redeye tetra), 37 *Carnegiella strigata* (marbled hatchetfish), 7 *Chilodus punctatus* (spotted headstander), and 240 *Astyanax bimaculatus* (twospot astyanax). The parasites found belonged to three taxa: monogeneans in the gills, nematodes (larvae of *Capillaria* sp. and *Contracaecum* sp.) in the digestive tract and liver and acanthocephalans (*Quadrigyrus torquatus*, *Q. brasiliensis* and *Q. nickoli*) in the stomach and intestine. *Astyanax bimaculatus* presented higher prevalence of acanthocephalans in the wet season, and lower prevalence of nematodes in the dry season. The possible importance of these parasites in the exportation of ornamental fish is discussed.

**Keywords:** Amazon, ornamental fish, parasite, acanthocephalans, monogeneans, nematodes.

## Resumo

O objetivo do presente trabalho foi avaliar a fauna parasitária de quatro espécies de peixes ornamentais capturados no rio Chumucuí, no município de Bragança-PA. Foram coletados um total de 307 peixes pertencentes a 4 espécies, sendo elas: *Moenkhausia sanctaefilomenae* (olho de fogo, n = 23), *Carnegiella strigata* (borboleta, n = 37), *Chilodus punctatus* (cabeça-para-baixo, n = 7) e *Astyanax bimaculatus* (lambari, n = 240) coletados de junho de 2006 a dezembro de 2007. Foram observados 3 taxa parasitando os peixes: monogenéticos nas brânquias, nematóides (larvas de *Capillaria* sp. e *Contracaecum* sp.) no trato digestório e fígado e acantocéfalos (*Quadrigyrus torquatus*, *Q. brasiliensis* e *Q. nickoli*) no estômago e intestino. *Astyanax bimaculatus* apresentou maior prevalência de acantocéfalos na estação chuvosa, menor prevalência de nematóides na estação seca. Discute-se a eventual importância destes parasitas na exportação de peixes ornamentais.

**Palavras-chave:** Amazônia, peixes ornamentais, parasita, acantocéfalos, monogenéticos, nematóides.

\*Corresponding author: Rodrigo Yudi Fujimoto  
Embrapa Tabuleiros Costeiros, Av. Beira Mar, 3250,  
CEP 49025-040, Aracajú, SE, Brasil  
e-mail: ryfujim@hotmail.com

## Introduction

The Amazon River basin is a hydrographical complex composed of a large number of rivers, lakes and small tributaries, containing great animal diversity. Within this area, fish are the most important resource for the local economy (MATOS et al., 2004).

The ornamental fishery is a major activity in that region and constitutes an important economic resource, since these fish are exported to other countries in large quantities (CHAO, 2001). The state of Pará is the second most important state regarding exportation of ornamental fish in Brazil, and the region of Bragança is an important place for catching fish because of the high number of different ornamental fish species. At least 38 different species have been reported in that region (Silva, unpublished data; Paula, unpublished data). This is especially evident in the Chumucuí River, a tributary of the Caeté River, where Characiform fish (which comprise a large number of ornamental species) represent 49% of the most important fish species (Silva, unpublished data).

It is well known that exportation of ornamental fish may represent a gateway for both ecto and endoparasites to enter a new environment. Their introduction may have serious effects, especially when parasites come into contact with new host fish. Several cases of introduced parasites, sometimes with devastating consequences, can be found in the literature (KIM et al., 2002; MORAVEC et al., 1999; EVANS; LESTER, 2001), and the pathology of parasites in ornamental fish has been described in a number of studies (FERRAZ; SOMMERVILLE, 1998; PIAZZA et al. 2006; THILAKARATNE et al., 2003; PRANG, 2007; MOUTON et al., 2001; GARCIA et al., 2009).

In this paper, ecto and endoparasites of some species of economically important ornamental fish species from the Chumucuí River were studied: *Carnegiella strigata* (Günther, 1864) (Gasteropelecidae), commonly named "marbled hatchetfish"; *Moenkhausia sanctaefilomenae* (Steindachner, 1907) (Characidae), "redeye tetra"; *Chilodus punctatus* (Müller & Troschel, 1844) (Chilodontidae), "spotted headstander"; and *Astyanax bimaculatus* (Linnaeus, 1758) (Characidae), commonly known as "two-spot lambari" or "matupiri" in the Amazon region, and as "piaba" in the north of the country. It is important to mention that the parasites of these fish species have never previously been studied in this region.

## Materials and Methods

Fish samples were collected twice a month (at the beginning and end of the month) from the Chumucuí River, a tributary of the Caeté River, in Bragança, Pará (1° 12' 38.3" S and 46° 47' 32" W).

Monthly sampling was done from June 2006 to December 2007. The fish were transported alive in Styrofoam boxes to the laboratory, where they were sacrificed using the anesthetic benzocaine. They were then measured (total and standard lengths) and weighed. Parasites were collected and processed as described by Eiras et al. (2006), and were identified at the lower taxonomic level in accordance with Thatcher (2006) and Travassos et al. (1928). The parasitological indexes were calculated as recommended by Bush et al. (1997).

The water pH and dissolved oxygen were recorded at the time of sampling. Environmental data (rainfall and temperature) were obtained from the Meteorological Station of Tracuateua, which belongs to the Brazilian National Institute of Meteorology.

The parasitological indexes were correlated with the weight, total length and standard length of the fish, and with the oxygen, pH, temperature and rainfall, by means of Pearson's correlation test. The variance of the mean values of prevalence and intensity of infection for the dry and wet seasons was analyzed and compared using the t test ( $P = 0.05$ ). Statistical tests were done using the Biostat 4.0 software.

## Results

Four species of ornamental fish were collected from the Chumucuí River: *Moenkhausia sanctaefilomenae*, *Carnegiella strigata*, *Chilodus punctatus* and *Astyanax bimaculatus*. The number of specimens, number of infected specimens, prevalence, total and standard lengths (cm) and total weight (g) are shown in Table 1. *Moenkhausia sanctaefilomenae* and *A. bimaculatus* presented the highest percentage prevalence; *C. punctatus* showed an intermediate percentage and *C. strigata* presented the lowest percentage prevalence, compared with the other species (Table 1).

The observations on the parasites showed that they belonged to three different taxa: nematodes (larvae of *Contracaecum* sp. and *Capillaria* sp.); acanthocephalans (*Quadrigyrrus torquatus*, Van Cleave, 1920; *Q. nickoli*, Schmidt and Huggins, 1973; *Q. brasiliensis*, Machado, 1941); and unidentified monogeneans. The distribution of the parasites according to the hosts is indicated in Table 2.

The infection by monogeneans was not related to the length of the host. However, *Moenkhausia sanctaefilomenae* showed the highest values for prevalence and mean intensity of infection by monogeneans (17.3% and  $5.0 \pm 2.2$ , respectively). *Astyanax bimaculatus* presented lower prevalence (2.08%) and similar mean intensity of infection ( $5.3 \pm 0.6$ ), compared with *M. sanctaefilomenae*. On the other hand, *C. strigata* showed lower prevalence (2.7%)

**Table 1.** Number of fish collected from Chumucuí River, number of infected fish, prevalence (%) of the infection, total length in cm, standard length in cm, and total weight in g.

|                                     | Fish collected | Infected fish | Prevalence (%) | Total length (cm) | Standard length (cm) | Total weight (g) |
|-------------------------------------|----------------|---------------|----------------|-------------------|----------------------|------------------|
| <i>Carnegiella strigata</i>         | 37             | 9             | 24.3           | $3.5 \pm 0.53$    | $2.8 \pm 0.36$       | $0.56 \pm 0.31$  |
| <i>Moenkhausia sanctaefilomenae</i> | 23             | 20            | 86.9           | $4.6 \pm 1.4$     | $3.9 \pm 1.03$       | $2.37 \pm 1.9$   |
| <i>Chilodus punctatus</i>           | 7              | 5             | 71.4           | $3.6 \pm 1.18$    | $4.8 \pm 1.06$       | $2.62 \pm 1.49$  |
| <i>Astyanax bimaculatus</i>         | 240            | 199           | 82.9           | $4.4 \pm 1.06$    | $3.6 \pm 0.95$       | $1.66 \pm 1.47$  |

and lower intensity of infection ( $1.0 \pm 0.16$ ). Monogeneans were not observed in *C. punctatus*.

Nematodes were found only as larval stages in different organs of the fish (Tables 3 and 4). *Astyanax bimaculatus* was the most parasitized species, and the highest level of nematode prevalence (50%) was in the stomach. *Chilodus punctatus* showed the highest mean intensity among the four fish species ( $12 \pm 0.3$ ). Larvae of *Contracaecum* sp. and *Capillaria* sp. were only found parasitizing the intestine and liver of *M. sanctaefilomenae*.

Concerning acanthocephalans, only cystacanth stages of *Quadrigyrus* spp. were found in the stomach and intestine.

**Table 2.** Distribution of the parasites according to fish species.

| Fish species               | Parasites  |
|----------------------------|--|
| <i>A. bimaculatus</i>      | <i>Q. torquatus</i> , <i>Q. nickoli</i> , unidentified monogeneans and nematodes   |
| <i>M. sanctaefilomenae</i> | <i>Contracaecum</i> sp., <i>Capillaria</i> sp. <i>Q. torquatus</i> , <i>Q. nickoli</i> , <i>Q. brasiliensis</i> and unidentified monogeneans |
| <i>C. punctatus</i>        | Unidentified nematodes and acanthocephalans  |
| <i>C. strigata</i>         | Unidentified monogeneans and nematodes   |

Only two host species presented larvae of *Quadrigyrus* sp. in the stomach: *C. punctatus* (prevalence 25%; mean intensity of infection  $1.5 \pm 0.74$ ) and *A. bimaculatus* (10.8%;  $1.76 \pm 0.63$ ). *Q. torquatus* and *Q. nickoli* were observed in *A. bimaculatus*; *Q. brasiliensis*, *Q. torquatus* and *Q. nickoli* were observed in *M. sanctaefilomenae*; and unidentified acanthocephalans were seen to be parasitizing *C. punctatus*.

The relationship between abiotic and biotic factors (Figure 1) showed that the only significant correlation with infection due to acanthocephalans was in the anterior intestine of *A. bimaculatus*, with higher prevalence in the wet season ( $P < 0.05$ ). In the same host, the prevalence of nematodes in the anterior intestine was also significantly higher in the dry season ( $P < 0.05$ ).

## Discussion


The results showed that parasites belonging to at least to three different taxa were present in the ornamental fish caught from the Chumucuí River. Similar results, sometimes with higher diversity of parasites, have been found in other ornamental fish species (PIAZZA et al., 2006; TAVARES-DIAS et al., 2009, 2010).

**Table 3.** Prevalence (%) of nematodes in the organs of the different fish species. Stomach; Anterior intestine; Middle intestine; Posterior intestine; Liver; Pyloric caecae; Total prevalence.

|                            | Stomach | Anterior intestine | Middle intestine | Posterior intestine | Liver | Pyloric caecae | Total prevalence (%) |
|----------------------------|---------|--------------------|------------------|---------------------|-------|----------------|----------------------|
| <i>A. bimaculatus</i>      | 49.1    | 27.91              | 12.0             | 34.7                | 36.25 | 9.1            | 82.9                 |
| <i>M. sanctaefilomenae</i> | 39.13   | 34.78              | 13.0             | 34.78               | 34.7  | 27.7           | 86.9                 |
| <i>C. punctatus</i>        | 50      | 37.5               | 0                | 1                   | 50    | 0              | 71.9                 |
| <i>C. strigata</i>         | 5.4     | 13.5               | 0                | 0                   | 2.7   | 0              | 24.3                 |

**Table 4.** Mean intensity (parasite/fish infected) of infection and standard deviation according to the nematodes in the organs of the different fish species. Stomach; Anterior intestine; Medium intestine; Posterior intestine; Liver; Pyloric caecae.

|                            | Stomach         | Anterior intestine | Middle intestine | Posterior intestine | Liver           | Pyloric caecae |
|----------------------------|-----------------|--------------------|------------------|---------------------|-----------------|----------------|
| <i>A. bimaculatus</i>      | $6.7 \pm 7.4$   | $4.49 \pm 3.04$    | $4 \pm 1.9$      | $10.41 \pm 1.83$    | $9.17 \pm 7.89$ | $3.1 \pm 1.3$  |
| <i>M. sanctaefilomenae</i> | $6.44 \pm 4.91$ | $3.55 \pm 2.6$     | $2 \pm 0.75$     | $4.5 \pm 3.69$      | $4.5 \pm 3.38$  | $8 \pm 4.7$    |
| <i>C. punctatus</i>        | $5.75 \pm 3.13$ | $5.6 \pm 3.8$      | 0                | $12 \pm 0.3$        | $10.7 \pm 3.1$  | 0              |
| <i>C. strigata</i>         | $2 \pm 0.4$     | $1.2 \pm 0.4$      | 0                | 0                   | $4 \pm 0.6$     | 0              |



**Figure 1.** Prevalence of parasites in the anterior intestine of the fish *Astyanax bimaculatus*, according to dry and wet seasons. a) nematode; b) acanthocephalan.

*Moenkhausia sanctaefilomenae* is a new host for *Contraecaecum* sp., *Capillaria* sp., *Q. torquatus*, *Q. nickoli* and *Q. brasiliensis*, and *A. bimaculatus* is a new host for *Q. torquatus* and *Q. nickoli*.

The monogeneans are mostly ectoparasites and are frequently observed in the gills. They are especially important in relation to fish rearing and may cause high mortality rates among farmed fish (FUJIMOTO et al., 2010). In these specimens, the intensity of infection was not related to the size of the hosts, contrary to what was observed by Pereira Junior et al. (2002) and Gonzáles et al. (2001) in relation to *Neoheterobothrium* parasitizing *Hippoglossina macrops*, which was related to greater surface area of the gills of the older hosts.

Larvae of the nematode species, *Contraecaecum* sp. and *Capillaria* sp., were observed parasitizing *M. sanctaefilomenae*. According to Dick and Choudhury (1995), the larvae may be more dangerous than the adults given that they can migrate through the fish to several target organs. *Astyanax bimaculatus* was the most infected fish species, presenting prevalence of 82%. It is interesting to note that for the same host species captured at Lages (state of Rio de Janeiro), Paraguassú and Luque (2007) reported prevalence of 41% for two nematode species, and in *A. fasciatus* they found prevalence of 39.2% for three nematode species. Also, Martins et al. (2005) found *Contraecaecum* sp. infecting 100% of *Hoplias malabaricus* and 80% of *Hoplyerythrinus unitaeniatus* caught in the state of Maranhão, and Barros et al. (2007) observed prevalence of 73% in *H. malabaricus* caught in the state of Mato Grosso. In the present study, the prevalence of nematodes in the anterior intestine was significantly higher in the dry season. However, the results of Martins et al. (2002) showed greater numbers of nematodes in the wet season in the Volta Grande region, state of Minas Gerais. According to Yamamoto et al. (2004), the alternation of wet and dry seasons and the consequent pronounced variation in the water level of the river could influence the food resources of the fish and indirectly influence the number of nematodes, due to variable predation on their intermediate hosts.

The acanthocephalans (*Q. torquatus*, *Q. nickoli* and *Q. brasiliensis*) were found in *A. bimaculatus*, *M. sanctaefilomenae* and *C. punctatus* hosts in the form of encysted larvae, while they were not observed in *C. strigata*. The presence in these hosts is probably related to predation on the intermediate hosts of the parasites, in which the feeding habits play an important role in the lifecycle. The related species *Q. machadoi* was reported in *Hemisorubim platyrhynchus* in the Paraná River (GUIDELLI et al., 2003), and in *H. malabaricus* in a lagoon at Aguai, infecting 87.5% of the fish (ROSIM et al., 2005). According to Rosim et al. (2005), *Astyanax altiparanae* is considered to be a paratenic host for *Q. torquatus*, and according to Carvalho et al. (2003), *Astyanax* spp. are considered both paratenic and intermediate hosts for *Q. torquatus*.

The results from these authors demonstrated that there was a negative correlation between the mean intensity of infection and the standard length of the host. Our observations do not support such evidence. The acanthocephalans were more prevalent in the anterior intestine of *A. bimaculatus* during the wet season, than the prevalence in the other hosts. The difference between hosts is probably due to different predation intensities of the intermediate hosts of the parasite, and the higher prevalence in *A. bimaculatus* during the wet season might be related to more abundant intermediate hosts during the wet season.

The fish of the present report are ornamental fish that are exported in great numbers to other countries. The results showed that these fish can carry several kind of parasites, both ecto and endoparasites, which can infect other fish in different continents and locations. In the literature, there are a number of reports dealing with the parasites of ornamental fish (FERRAZ; SOMMERVILLE, 1998; PIAZZA et al., 2006; THILAKARATNE et al., 2003; PRANG, 2007; MOUTON et al., 2001; GARCIA et al., 2009) and with the introduction of parasites by ornamental fish (KIM et al., 2002; MORAVEC et al., 1999; EVANS; LESTER, 2001). The consequences of parasite introduction can be detrimental to native fish, sometimes causing serious epizootics that may lead to extensive mortality as shown in relation to several species of monogeneans introduced into aquariums (BULLARD et al., 2001), into contact with farmed fish (OGAWA et al., 1995; STERUD et al., 1998) and also into natural populations with dramatic consequences (WHITTINGTON; CHISHOLM, 2003; BAKKE et al., 2002).

The parasites observed might constitute a threat to other fish species in the region where they have been introduced. Therefore, it is recommended that prophylactic measures should be taken prior to exportation of fish, and that fish should undergo a period of quarantine at the place of introduction. In the present sample, three groups of endoparasites with heteroxenous life cycles (nematodes and acanthocephalans) were found and one group of ectoparasites with a direct life cycle (monogeneans). Consequently, the prophylactic measures must be especially directed towards monogeneans, for several reasons: they are located in the gills, a particularly fragile and important fish organ; they reproduce very quickly in a closed environment given that they have a direct life cycle; and because they are ectoparasites, use of appropriate drugs is easier and more effective.

## Acknowledgements

The participation of J.C. Eiras on this research was partially supported by the European Regional Development Fund (ERDF) through the COMPETE - Operational Competitiveness Programme and national funds through FCT - Foundation for Science and Technology, under the project "PEst-C/MAR/LA0015/2011".

## References

- Bakke TA, Harris PD, Cable J. Host specificity dynamics: observations on gyrodactylid monogeneans. *Int J Parasitol* 2002; 32(3): 281-308. [http://dx.doi.org/10.1016/S0020-7519\(01\)00331-9](http://dx.doi.org/10.1016/S0020-7519(01)00331-9)
- Barros LA, Moraes Filho J, Oliveira RL. Larvas de nematóides de importância zoonótica encontradas em traíras (*Hoplias malabaricus* bloch, 1794) no município de Santo Antonio do Leverger, MT. *Arq Bra Med Vet Zootec* 2007; 59(2): 533-535. <http://dx.doi.org/10.1590/S0102-09352007000200042>
- Bullard SA, Frasca SJR, Benz GW. Gill lesions associated with *Erpocotyle tiburonis* (Monogenea: Hexabothriidae) on wild and aquarium-held bonnet head sharks (*Sphyrna tiburo*). *J Parasitol* 2001; 87: 972-977. PMID:11695418.
- Bush AO, Lafferty KD, Lotz JM, Shostak AW. Parasitology meets ecology on its own terms: Margolis et al revisited. *J Parasitol* 1997; 83(4): 575-583. PMID:9267395. <http://dx.doi.org/10.2307/3284227>

- Carvalho S, Guidelli GM, Takemoto RM, Pavanelli GC. Ecological aspects of endoparasite fauna of *Acestrorhynchus lacustris* (Lütken, 1875) (Characiformes, Acestrorhynchidae) on the Upper Paraná River floodplain, Brazil. *Acta Sci Biol Sci* 2003; 25(2): 479-483. <http://dx.doi.org/10.4025/actasciobiolsci.v25i2.2043>
- Chao NL. The fishery diversity and conservation of ornamental fishes in the Rio Negro Basin, Brasil: A review of Project Piaba (1989-1999). In: Chao NL, Petry P, Prang G, Sonnenschein L, Tlusty MT, editors. *Conservation and management of ornamental fish resources of the Rio Negro Basin, Amazonia Brasil - Project Piaba*. Manaus: Universidade do Amazonas, Manaus; 2001. p. 161-204.
- Dick TA, Choudhury A. Phylum nematoda. In: Woo PTK, editors. *Fish diseases and Disorders*. New York: CABE Publishing; 1995. v. 1. Protozoan and metazoan infections, p. 415-446.
- Eiras JC, Takemoto RM, Pavanelli GC. *Métodos de estudo e técnicas laboratoriais em parasitologia de peixes*. 2. ed. Maringá: Eduem; 2006.
- Evans BB, Lester RJG. Parasites of ornamental fish imported into Australia. *EAFP Bul* 2001; 21(2): 51-55.
- Ferraz E, Sommerville C. Pathology of *Piscinoodinium* sp. (Protozoa: Dinoflagellida), parasites of the ornamental freshwater catfishes *Corydoras* spp. and *Brochis splendens* (Pisces: Callichthyidae). *Dis Aquat Org* 1998; 33(1): 43-49. PMID:9653457. <http://dx.doi.org/10.3354/dao033043>
- Fujimoto RY, Castro MP, Martins ML, Moraes FR, Varella JEA, Diniz DG. Effects of Chromium Supplementation on the Infrapopulations of *Anacanthorus penilabiatus* (Monogeneoidea) and *Piscinoodinium pillulare* (Dinoflagellida) Parasites of *Piaractus mesopotamicus* (Characidae). *Braz Arch Biol Technol* 2010; 53(4): 827-833. <http://dx.doi.org/10.1590/S1516-89132010000400011>
- Garcia F, Fujimoto RY, Martins ML, Moraes FR. Protozoan parasites of *Xiphophorus* spp. (Poeciliidae) and their relation with water characteristics. *Arq Bra Méd Vet Zootec* 2009; 61(1): 156-162. <http://dx.doi.org/10.1590/S0102-09352009000100022>
- Guidelli GM, Isaac A, Takemoto RM, Pavanelli GC. Endoparasite infracommunities of *Hemisorubim platyrhynchos* (Valenciennes, 1840) (Pisces: Pimelodidae) of the Baía River, upper Paraná River floodplain, Brazil: specific composition and ecological aspects. *Braz J Biol* 2003; 63(2): 261-268. PMID:14509848. <http://dx.doi.org/10.1590/S1519-69842003000200011>
- González MT, Acuña E, Oliva ME. Metazoan parasite fauna of the Bigeye Flounder, *Hippoglossina macrops*, from Northern Chile. Influence of host age and sex. *Mem Inst Oswaldo Cruz* 2001; 96(8): 1049-1054.
- Kim JH, Hayward CJ, Joh SJ, Heo GJ. Parasitic infections in live freshwater tropical fishes imported to Korea. *Dis Aquat Org* 2002; 52(2): 169-173. PMID:12542094. <http://dx.doi.org/10.3354/dao052169>
- Martins ML, Mello A, Paiva FC, Fujimoto RY, Schalch SHC, Colombano NC. Prevalência, sazonalidade e intensidade de infecção por *Diplostomum* (*Austrodiplostomum*) *compactum* Lutz, 1928 (Digenea, Diplostomidae), em peixes do reservatório de Volta Grande, Estado de Minas Gerais, Brasil. *Acta Sci* 2002; 24(2): 469-474.
- Martins ML, Onaka EM, Fenerick Junior J. Larval *Contracaecum* sp. (Nematoda: Anisakidae) in *Hoplias malabaricus* and *Hoplerhynchus unitaeniatus* (Osteichthyes: Erythrinidae) of economic importance in occidental marshlands of Maranhão, Brazil. *Vet Parasitol* 2005; 127(1): 51-59. PMID:15619375. <http://dx.doi.org/10.1016/j.vetpar.2004.09.026>
- Matos E, Casal G, Matos P, Corral LA, Carlos. Microorganismos Parasitos de Animais Aquáticos da Amazônia. In: Ranzani-Paiva MJT, Takemoto RM, Lizama MAP, editors. *Saúde de organismos aquáticos*. Varella; 2004. p.159-178.
- Moravec F, Wolter J, Körting W. Some nematodes and acanthocephalans from exotic ornamental freshwater fishes imported into Germany. *Folia Parasitol* 1999; 46(4): 296-310.
- Mouton A, Basson L, Impson D. Health status of ornamental freshwater fishes imported to South Africa: a pilot study. *Aquar Sci Conserv* 2001; 3(4): 313-319. <http://dx.doi.org/10.1023/A:1013197928590>
- Ogawa K, Bondad-Reantaso MG, Fukudome M, Wakabayashi H. *Neobenedenia girellae* (Hargis, 1955) Yamaguti, 1963 (Monogenea: Capsalidae) from cultured marine fishes of Japan. *J Parasitol* 1995; 81(2): 223-227. PMID:7707197. <http://dx.doi.org/10.2307/3283923>
- Paraguassú AR, Luque JL. Metazoários Parasitos de seis espécies de Peixes do Reservatório de Lajes, Estado do Rio de Janeiro, Brasil. *Rev Bras Parasitol Vet* 2007; 16(3): 121-128. PMID:18078597. <http://dx.doi.org/10.1590/S1984-29612007000300002>
- Pereira Junior J, Da Costa MAS, Vianna RT. Índices parasitológicos de Cucullanidae (Nematoda: Seuratoidea) em *Micropogonias furnieri* (Desmarest, 1823) no litoral do Rio Grande do Sul, Brasil. *Atlântica* 2002; 24(2): 97-101.
- Piazza RS, Martins ML, Guiraldelli L, Yamashita M. Parasitic diseases of freshwater ornamental fishes commercialized in Florianópolis, Santa Catarina, Brazil. *B Inst Pesca* 2006; 32(1): 51-57.
- Prang G. An industry analysis of the freshwater ornamental fishery with particular reference to the supply of Brazilian freshwater ornamentals to the UK market. *Uakari* 2007; 3(1): 7-51.
- Rosim DF, Cecarelli PS, Silva-Souza AT. Parasitismo de *Hoplias malabaricus* (Bloch, 1794) (Characiformes, Erythrinidae) por *Quadrigrayrus machadoi* Fábio, 1983 (Eoacanthocephala, Quadrigrayridae) de uma Lagoa em Aguai, Estado De São Paulo, Brasil. *Rev Bras Parasitol Vet* 2005; 14(4): 147-153. PMID:16445871.
- Sterud E, Harris PH, Bake TA. The influence of *Gyrodactylus salaris* Malmberg 1957 (Monogenea) on the epidermis of Atlantic salmon, *Salmo salar* L., and brook trout, *Salvelinus fontinalis* (Mitchill): experimental studies. *J Fish Dis* 1998; 21(4): 257-263. <http://dx.doi.org/10.1046/j.1365-2761.1998.00099.x>
- Tavares-Dias M, Lemos JRG, Martins ML, Jerônimo GT. Metazoan and protozoan parasites of freshwater ornamental fish from Brazil. In: Tavares-Dias M, editors. *Manejo e sanidade de peixes em cultivo*. Macapá: Embrapa; 2009. p. 469-494.
- Tavares-Dias M, Gonzaga Lemos JR, Martins ML. Parasitic fauna of eight species of ornamental freshwater fish species from the middle Negro River in the Brazilian Amazon Region. *Rev Bras Parasitol Vet* 2010; 19(2): 103-107. PMID:20624347. <http://dx.doi.org/10.4322/rbpv.01902007>
- Thatcher VE. *Amazon Fish Parasites*. Manaus: Instituto Nacional de pesquisas da Amazônia; 2006. p. 555.
- Thilakarathne IDSIP, Rajapaksha G, Hewakopara A, Rajapakse RPVJ, Faizal ACM. Parasitic infections in freshwater ornamental fish in Sri Lanka. *Dis Aquatic Org* 2003; 54(2): 157-162. PMID:12747641. <http://dx.doi.org/10.3354/dao054157>
- Travassos L, Artigas P, Pereira C. Fauna helmintológica dos peixes de água doce do Brasil. *Arch Inst Biol* 1928; 1: 5-68.
- Whittington ID, Chisholm LA. Biodiversity of marine parasites in Australia; more than just a list of largely invisible creatures. *Rec South Aust Mus* 2003; 7: 51-60.
- Yamamoto KC, Soares MGM, Freitas CEC. Alimentação de *Triporthesus angulatus* (Spix & Agassiz, 1829) no lago Camaleão, Manaus, AM, Brasil. *Acta Amaz* 2004; 34(4): 653-659. <http://dx.doi.org/10.1590/S0044-59672004000400017>