

Whole-Genome Sequence of *Corynebacterium pseudotuberculosis* Strain Cp162, Isolated from Camel

Syed Shah Hassan,^a Maria Paula C. Schneider,^b Rommel Thiago Jucá Ramos,^b Adriana R. Carneiro,^b Alex Ranieri,^b Luis Carlos Guimarães,^b Amjad Ali,^a Syeda Marriam Bakhtiar,^a Ulisses de Pádua Pereira,^e Anderson Rodrigues dos Santos,^a Siomar de Castro Soares,^a Fernanda Dorella,^a Anne Cybelle Pinto,^a Dayana Ribeiro,^a Maria Silvanira Barbosa,^b SÍntia Almeida,^a Vinícius Abreu,^a Flávia Aburjaile,^a Karina Fiaux,^a Eudes Barbosa,^a Carlos Diniz,^a Flavia S. Rocha,^a Rashmi Saxena,^c Sandeep Tiwari,^{c,g} Vasudeo Zambare,^c Preetam Ghosh,^{c,d} Luis G. C. Pacheco,^h Christopher G. Dowson,^f Anil Kumar,^g Debmalya Barh,^c Anderson Miyoshi,^a Vasco Azevedo,^a and Artur Silva^b

Instituto de Ciências Biológicas, Universidade Federal de Minas Gerais, Belo Horizonte, MG, Brazil^a; Instituto de Ciências Biológicas, Universidade Federal do Pará, Belém, PA, Brazil^b; Centre for Genomics and Applied Gene Technology, Institute of Integrative Omics and Applied Biotechnology (IIOAB), Nonakuri, Purba Medinipur, West Bengal, India^c; Department of Computer Science and Center for the Study of Biological Complexity, Virginia Commonwealth University, Richmond, Virginia, USA^d; Departamento de Medicina Veterinária, Universidade Federal de Lavras, Lavras, Brazil^e; School of Life Sciences, University of Warwick, Coventry, United Kingdom^f; School of Biotechnology, Devi Ahilya University, Indore, India^g; and Instituto de Ciências da Saúde, Universidade Federal da Bahia, Salvador, Brazil^h

***Corynebacterium pseudotuberculosis* is a pathogen of great veterinary and economic importance, since it affects livestock, mainly sheep and goats, worldwide, together with reports of its presence in camels in several Arabic, Asiatic, and East and West African countries, as well as Australia. In this article, we report the genome sequence of *Corynebacterium pseudotuberculosis* strain Cp162, collected from the external neck abscess of a camel in the United Kingdom.**

Corynebacterium pseudotuberculosis is a facultative intracellular pathogen which causes caseous lymphadenitis (CLA) in sheep and goats, ulcerative lymphangitis in horses, and diseases in cattle and in humans (11). In camels, *C. pseudotuberculosis* affects almost 10% of the population in a herd (1). Arthritis, peri-arthritis, and pleuritis in camels have been found in Gran Canaria island, Spain (13).

CLA causes the formation of abscesses in large superficial lymph nodes and occasionally in the lymph nodes of internal organs in small ruminants and mammals other than humans, yet it has a different lesion nature in camels (5). Early reports of CLA in sheep from Saudi Arabia (8) and nearby Middle East countries describe the disease prevalence in camel herds. The most prevalent biovar in camelids has long been described as biovar ovis (15), but further studies have indicated the susceptibility of dromedary camels to biovar equi in a close environment with CLA-affected horses or cattle (14).

CLA was first reported for United Kingdom goat and sheep flocks in 1990 and 1991, respectively (4, 10). The *C. pseudotuberculosis* strain Cp162 was obtained from an external neck abscess of a camel in 1999 and was originally designated W00644. The biochemical characteristics were determined using the API Coryne test (bioMérieux, Marcy-l'Étoile, France), according to the manufacturer's instructions. Cp162 is nitrate positive and was typed using a multilocus sequence typing (MLST) technique by the Moredun Research Institute whose sequence type (ST) was 3 (2).

We sequenced the genome of Cp162 using a fragment library of the SOLiD v3 Plus sequencing platform (Applied Biosystems), generating 75,659,346 reads of 50 bp in length. After quality filtering (9), reads representing an average Phred quality of less than 20 were removed, followed by the correction of sequencing errors with the SAET software program. In total, 31,568,756 reads with a high quality score were selected, with 686-fold coverage, considering the size of the reference genome used (2.3 Mb), that of *C. pseudotuberculosis* strain FRC41 (NC_014329). The reads were mapped against the reference genome of *C. pseudotuberculosis* 316

(CP003077) through the CLC Genomics Workbench, generating a genome scaffold containing gaps, which were resolved by recursive alignments of short reads (3) and manual curation.

The functional annotation of the genome for open reading frame (ORF) prediction was performed using the software program FgenesB (Softberry). The software program RNAmmer (6) was used for the prediction of rRNA using hidden Markov models (HMM); to identify transfer RNAs (tRNAs), the tRNAscan-SE program was used (7).

Identification of protein domains and families was performed with InterProScan software (16), which integrates several related databases. The nonredundant protein database of NCBI (nr) was used to characterize the coding sequences (CDS) with the help of the Artemis software tool (12). Thus, the genome of Cp162, having a size of 2,293,464 bp, 52.17% GC content, 2,002 coding sequences, 4 rRNA operons, 49 tRNA genes, and 87 pseudogenes, was deposited in the NCBI database.

Further genome analysis will provide deep insight into the virulence of this pathogen at the molecular and genetic levels and will aid in detecting effective vaccine and drug targets in postgenomic studies.

Nucleotide sequence accession number. The genome sequence obtained in this study has been deposited in the GenBank database under accession number [CP003652](https://doi.org/10.1128/JB.01373-12).

ACKNOWLEDGMENTS

This work is the outcome of a collaboration between various prestigious organizations, with the support of longstanding research and academic institutions, including the Rede Paraense de Genômica e Proteômica,

Received 30 July 2012 Accepted 6 August 2012

Address correspondence to Artur Silva, asilva@ufpa.br.

Copyright © 2012, American Society for Microbiology. All Rights Reserved.

doi:10.1128/JB.01373-12

supported by the Fundação de Amparo a Pesquisa do Estado do Pará, the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), and the Fundação de Amparo a Pesquisa do Estado de Minas Gerais (FAPEMIG). M.P.C.S., V.A., and A.S. were supported by the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq). We also acknowledge support from the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES).

REFERENCES

1. Afzal M, Sakir M, Hussain MM. 1996. *Corynebacterium pseudotuberculosis* infection and lymphadenitis (taloa or mala) in the camel. *Trop. Anim. Health Prod.* **28**:158–162.
2. Bolt F. 2009. The population structure of the *Corynebacterium diphtheriae* group. Ph.D. dissertation. University of Warwick, Coventry, United Kingdom.
3. Cerdeira LT, et al. 2011. Rapid hybrid de novo assembly of a microbial genome using only short reads: *Corynebacterium pseudotuberculosis* I19 as a case study. *J. Microbiol. Methods* **86**:218–223.
4. Harker DB. 1990. Investigation of an outbreak of caseous lymphadenitis in goats in the United Kingdom. *Goat Vet. Soc. J.* **11**:52–54.
5. Hawari AD. 2008. *Corynebacterium pseudotuberculosis* infection (caseous lymphadenitis) in camels (*Camelus dromedarius*) in Jordan. *Am. J. Anim. Vet. Sci.* **3**:68–72.
6. Lagesen K, et al. 2007. RNAmmer: consistent and rapid annotation of ribosomal RNA genes. *Nucleic Acids Res.* **35**:3100–3108.
7. Lowe TM, Eddy SR. 1997. tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res.* **25**:955–964.
8. Radwan AI, el-Magawry S, Hawari A, al-Bekairi SI, Rebleza RM. 1989. *Corynebacterium pseudotuberculosis* infection in camels (*Camelus dromedarius*) in Saudi Arabia. *Trop. Anim. Health Prod.* **21**:229–230.
9. Ramos RT, et al. 2011. Analysis of quality raw data of second generation sequencers with Quality assessment Software. *BMC Res. Notes* **4**:130. doi:10.1186/1756-0500-4-130.
10. Robins R. 1991. Focus on—caseous lymphadenitis. *State Vet. J.* **1**:7–10.
11. Ruiz JC, et al. 2011. Evidence for reductive genome evolution and lateral acquisition of virulence functions in two *Corynebacterium pseudotuberculosis* strains. *PLoS One* **6**:e18551. doi:10.1371/journal.pone.0018551.
12. Rutherford K, et al. 2000. Artemis: sequence visualization and annotation. *Bioinformatics* **16**:944–945.
13. Tejedor-Junco MT, Lupiola P, Caballero MJ, Corbera JA, Gutierrez C. 2009. Multiple abscesses caused by *Salmonella enterica* and *Corynebacterium pseudotuberculosis* in a dromedary camel. *Trop. Anim. Health Prod.* **41**:711–714.
14. Tejedor-Junco MT, Lupiola P, Schulz U, Gutierrez C. 2008. Isolation of nitrate-reductase positive *Corynebacterium pseudotuberculosis* from dromedary camels. *Trop. Anim. Health Prod.* **40**:165–167.
15. Tejedor MT, Martin JL, Corbera JA, Shulz U, Gutierrez C. 2004. Pseudotuberculosis in dromedary camels in the Canary Islands. *Trop. Anim. Health Prod.* **36**:459–462.
16. Zdobnov EM, Apweiler R. 2001. InterProScan—an integration platform for the signature-recognition methods in InterPro. *Bioinformatics* **17**:847–848.