



Comparative cytogenetic analysis in the species *Uroderma magnirostrum* and *U. bilobatum* (cytotype $2n = 42$) (Phyllostomidae, Stenodermatinae) in the Brazilian Amazon

Adailton Moreira da Silva¹, Suely Aparecida Marques-Aguiar², Regina Maria de Souza Barros³, Cleusa Yoshiko Nagamachi³ and Julio Cesar Pieczarka³

¹Universidade Estadual do Amazonas, Centro de Estudos Superiores de Parintins, Parintins, AM, Brazil.

²Museu Paraense Emílio Goeldi, Departamento de Zoologia, Belém, PA, Brazil.

³Universidade Federal do Pará, Departamento de Genética, Centro de Ciências Biológicas, Laboratório de Citogenética, Belém, PA, Brazil.

Abstract

The genus *Uroderma* includes two species: *U. magnirostrum* and *U. bilobatum*. These species are characterized by their high degree of karyotypic evolution, diverging from most other species of the subfamily Stenodermatinae, which have a lower degree of chromosomal evolution. The present study reports the first banding patterns of *U. magnirostrum* (G-, C-banding and Ag-NOR) and *U. bilobatum* (C-banding and Ag-NOR). The chromosomal data in conventional staining of *U. magnirostrum* ($2n = 36$, NF = 62) and *U. bilobatum* (cytotype $2n = 42$, NF = 50) are equivalent to that described in the literature. When compared, chromosomal homeologies are found in both karyotypes, as well as differences, confirming that karyotypic evolution in the *Uroderma* genus is intense. Fission, fusion, inversion or translocation events are required to explain the karyotypic evolution of this genus. The comparison of karyotype, described here, to one of the species of the genus *Artibeus* ($2n = 30/31$), suggests that some chromosomal forms are apomorphic and shared between the two species of *Uroderma*. This confirms the monophyly of the genus, and that *U. magnirostrum* presents a more primitive karyotype when compared to *U. bilobatum*.

Key words: cytogenetics, chromosomal evolution, Phyllostomidae, *Uroderma*.

Received: November 16, 2004; Accepted: December 14, 2004.

Introduction

The *Uroderma* genus is a group of Stenodermatines bats primarily recognized as frugivores with slightly round faces, small snout and lancet nose. They also possess four facial stripes and a very evident dorsal stripe (Nowak, 1996). *Uroderma* comprises two species, *U. bilobatum* (common) and *U. magnirostrum* (uncommon), and their geographic range encompasses southern Mexico to southeastern Brazil (Hill and Smith, 1984; Nowak, 1996).

The Stenodermatinae subfamily comprises 18 genera (Nowak, 1996) and the relationships of the *Uroderma* genus with other members of the subfamily are the subject of discussion. The phylogenetic proposals using data from different characters show different associations of the genus *Uroderma* with other genera of Stenodermatinae.

Using karyotypic information, Gardner (1977) placed this genus as a different lineage, occupying a basal position in the subfamily. Using morphological information, Owen (1987) related *Uroderma* with *Ectophylla* and *Artibeus* genera. Using molecular data, Van Den Bussche (1992) suggested that *Uroderma* is more closely related to *Vampyrops* and *Vampyrodes* than with the other genera. With the help of morphological and karyotypic information, Lim (1993) related this genus strictly to *Artibeus*, in spite of the great chromosomal divergence among them.

This genus is an exception to the karyotypic conservation found in most species of the Stenodermatinae subfamily, because it presents both intraspecific and interspecific variability (Baker, 1967, 1973, 1979; Varela-Garcia *et al.*, 1989). The species *U. magnirostrum* had its karyotype described by Baker and Lopez (1970), without banding information. *U. bilobatum* presents intraspecific variability with, at least, three chromosomal cytotypes: $2n = 38$, 42 and 44 (Baker and Lopez, 1970;

Send correspondence to Adailton Moreira da Silva. Universidade Estadual do Amazonas, Centro de Estudos Superiores de Parintins, Estrada Odovaldo Novo s/n, 69152-470 Parintins, AM, Brazil. E-mail: amdsilva@uea.edu.br.

Baker *et al.*, 1972, 1975, 1982). The cytotypes with $2n = 38$ and $2n = 44$ are derived from three mutational events; two translocation and one centric fusion (Baker, 1979, 1981). The cytotypes with $2n = 42$ and $2n = 44$ are distinguished by one centric fusion and one pericentric inversion (Baker *et al.*, 1982).

The Amazonian area is, without a doubt, one of the largest centers of biodiversity on the planet, but little is known about the cytogenetics of the species that are part of this diversity. The objective of this work is to study the karyotype of *U. magnirostrum* and *U. bilobatum* (cytotype $2n = 42$) species, collected in the Brazilian Amazon using G-, C-banding and Ag-NOR staining, and to compare the data to the karyotypes already described for both species. They will also be compared to the karyotype of *Artibeus* ($2n = 30/31$), which is supposed to be the primitive karyotype of the subfamily. This information will be used to define the pattern of chromosomal evolution within the genus *Uroderma*.

Material and Methods

The sample consists of four *U. magnirostrum* (3M and 1F) and two *U. bilobatum* (1M and 1F) collected in Belém, Pará, Brazil and were deposited in the Mammal collection at the Museu Paraense Emílio Goeldi (MPEG 27271, 27272, 27476, 27557, 27568, 27572). The animals were collected according to Bredt *et al.* (1996), with the use of mist nets.

The metaphase chromosomes were obtained using the technique of direct extraction of bone marrow, according to Ford and Hamerton (1956). The animals were previously submitted to mitotic induction by yeast stimulation, according to Lee and Elder (1980). The G-bands were obtained by incubating slides in a solution of 2XSSC (60 °C) from 2 to 10 min and staining in a Wright Giemsa solution (0.25% added to 3 mL of phosphate buffer, pH 6.8) for 2 min and 30 s. The C-banding was performed according to Sumner (1972). The Nucleolar Organizer Regions (NORs) staining was performed according to Howell and Black (1980).

The chromosomes were grouped according to morphology and in decreasing size, following Baker and Lopez (1970).

Results

The *U. magnirostrum* specimens presented $2n = 36$, $NF = 62$ (Figure 1a). The autosome set is composed of 4 metacentric, 10 submetacentric and 3 acrocentric pairs. The X chromosome is a submetacentric of medium size and the Y is a small submetacentric. In one specimen the submetacentric pair 5 showed a heteromorphism in the short arm. The C-banding pattern (Figure 1b) shows that the constitutive heterochromatin is located in the pericentromeric region of all the autosomes and in the distal




Figure 1 - G- (a) and C-banded (b) karyotypes of *U. magnirostrum*. Note the heteromorphism of pair 5.

region of the short arm of all the submetacentrics. The few exceptions are the heteromorphic pair 5 whose homologues have a interstitial band in the short arm and the pair 6 that has proximal band in the long arm. The X chromosome has centromeric and terminal heterochromatin in the short arm. The Y has a heterochromatic block in the long arm. The number of Ag-NORs ranges from five to eight per metaphase, and they are located in the distal region of the short arm of submetacentric chromosomes (Figure 3a).

The *U. bilobatum* specimens had $2n = 42$, $NF = 50$ (Figure 2a). The autosome set is composed of 5 submeta-

centric and 15 acrocentric pairs. The X chromosome is a medium sized submetacentric and the Y, a small submetacentric. The C-banding pattern (Figure 2b) shows the constitutive heterochromatin in the pericentromeric region of all the autosomes and in the distal region of the short arm of two submetacentric chromosomes (pairs 1 and 2). The X chromosome shows centromeric and terminal heterochromatin in the short arm. The Y has a block of heterochromatin in the long arm. The Ag-NORs were located in the distal region of the short arm of submetacentric/acrocentric chromosomes (Figure 3b), ranging from five to eight per metaphase.




Figure 2 - G- (a) and C-banded (b) karyotypes of *Uroderma bilobatum*.

The G-banding comparison of the karyotypes of the two species of *Uroderma* (Figure 4) shows homeologies of whole chromosomes (Figure 4a), segments or arms of chromosomes (Figures 4b, 4c and 4d) and chromosomes that have no correspondence (Figures 4e and 4f). Both species present a chromosomal system of sexual determination of the type Neo-XY, but the G-banding obtained on the sex chromosomes, mainly the X, demonstrates that these chromosomes have no similar banding patterns (Figure 4g). The C-banded karyotype of *U. magnirostrum* (Figure 1b) shows a larger amount of heterochromatin than *U. bilobatum* (Figure 2b). The 5-8 multiple Ag-NORs were located in different chromosomes as can be seen in Figure 3.

Discussion

The *U. magnirostrum* species, collected in the Brazilian Amazon, has a similar karyotype ($2n = 36$, $NF = 62$) to that published by Baker and Lopez (1970) from animals collected in Central America, whose chromosomes were analyzed under giemsa staining. The authors also found two males with $2n = 35$, $NF = 62$, due to the presence of a large metacentric chromosome (possibly originating from a fusion of two acrocentrics), not seen in the present work.




Figure 3 - Metaphases of *U. magnirostrum* (a) and *U. bilobatum* (b) showing chromosomes with Ag-NORs (arrow).




Figure 4 - Karyotypic comparison between *Uroderma bilobatum* (UBI) and *U. magnirostrum* (UMA). a) Homeologies of whole chromosomes: 1, 2, 10, 12 and 15 of UBI, and 7, 9, 15, 16 and 17 of UMA; b) Homeologies of the short arms (p) of the UMA1 with the acrocentric pair UBI14, while the long arms (q) correspond to the acrocentric pair UBI13, with a paracentric inversion; c) The long arm of pair UMA8 is similar to proximal region of long arm of the pair UBI6, while terminal region of long arm of pair UBI6 is similar to the long arm of pair UMA2; d) Homeologies of the long arm of the pairs 5, 6 and 11 of UMA with the acrocentric pairs 7, 8 and 9 of UBI, respectively; e) Pairs of UBI that have no correspondence in the karyotype of UMA; f) Pairs of UMA that have no correspondence in the karyotype of UBI; g) Comparison of the sex chromosomes: both have chromosomal system of sexual determination of the type Neo-XY.

The unique variation observed was a specimen that showed a heteromorphism in the short arm of the largest submetacentric (chromosome 5). At least three hypotheses could be advanced to explain this heteromorphism: 1) a heterochromatinization in only one of the homologues (see C-banding); 2) amplification of the rDNA cistron (see Ag-NOR banding); 3) a deletion in the short arm of only one of the homologues.

The *U. bilobatum* species (cytotype $2n = 42$), has a similar karyotype to that published by Baker and Lopez (1970) and Baker *et al.* (1982). From a chromosomal evolution viewpoint, this species is very important, because it possesses karyotypes ranging from $2n = 38$ to 44 (Baker and Lopez, 1970; Baker 1979, 1981; Baker *et al.*, 1982). According to Baker *et al.* (1982), two evolutionary histories for these karyotypes can be inferred. In the first, there is a trichotomy, because the three karyotypes appeared at the same time, and the cytotype $2n = 44$ would represent the primitive condition. In the second, a fusion and an inversion separate a group with $2n = 42$ and other with $2n = 44/38$ and two translocations/fusion separate the $2n = 38$ from the $2n = 44$. However, even using the data analyzed here and those from the literature, the chromosomal phylogeny of the cytotypes of the *U. bilobatum* species remains uncertain, and more chromosomal information is necessary to solve this puzzle.

The high degree of karyotypic homeologies found between *Uroderma* species reinforces its relationship. They possess exclusive features such as pair 15 of *U. magnirostrum* or pair 10 of *U. bilobatum* not found in any other species of Stenodermatinae bats with G-banded karyotypes described so far. These species also shared two pairs of chromosome (16 and 17 of *U. magnirostrum*; 12 and 15 of *U. bilobatum*) that arose by fission of a metacentric chromosome. These chromosomes have some similarities to the chromosome 4 of *A. lituratus* (Silva, 2000) and 5 of *Phyllostomus hastatus* (Rodrigues, 1998). According to Baker *et al.* (1979), the karyotypes of the members of the Stenodermatinae subfamily have two chromosomes that are not found in any other species of the family Phyllostomidae, and are considered a plesiomorphic character in the *Uroderma* species.

Taking into account that the karyotype $2n = 30/31$ of *Artibeus* represents the most primitive for the subfamily (Baker *et al.*, 1979; Silva, 2000) and that *U. magnirostrum* might present the most primitive karyotype for the genus, because of the highest similarity to *Artibeus*, we can infer karyoevolutionary pathways for the genus *Uroderma*: 1) one paracentric inversion in the long arm and subsequent fission of the pair 1 of *U. magnirostrum* originated the pairs 13 and 14 of *U. bilobatum*; 2) at least, two fission events and one fusion *in tandem* is necessary to explain the homeologies of the chromosomal segments among pair 6 of *U. bilobatum* and pairs 2 and 8 of *U. magnirostrum*; 3) fission events might have occurred also, to give rise to the

acrocentric chromosomes 7, 8 and 9 of *U. bilobatum* from the submetacentric chromosomes 5, 6 and 11 of *U. magnirostrum*, respectively. For the other chromosomes, the resolution of G-bands was too low to allow good comparison in small euchromatic areas that suffered chromosomal rearrangements. However, new techniques of molecular cytogenetics (such as FISH) can help explain doubts that still remain on the chromosomal evolution of this genus.

Regarding the chromosomal systems of sex determination of these two species (neo-XY/neo-XX), Tucker (1986) and Tucker and Bickham (1986) proposed that two chromosomal rearrangements possibly gave rise to the sex chromosomes observed in the karyotypes of the species of Phyllostomidae. First, a translocation X-autosome gave rise to a chromosomal pattern XY_1Y_2 , as observed in most of the species of the *Artibeus* genus. Second, a translocation Y-autosome originated the Neo-XY pattern, as observed in the species of the present work. Thus, the chromosomal system of sex determination neo-XY is probably derived from system XY_1Y_2 , as corroborated by the similarity in the size and in the C-banding pattern of the acrocentric chromosomes of the system XY_1Y_2 (genus *Artibeus*) and metacentric chromosomes of the system neo-XY (genus *Uroderma*).

Our comparative analysis of the two species of *Uroderma* allowed us to conclude that they are monophyletic. The species *U. bilobatum* presents an apomorphic karyotype, while whole *U. magnirostrum* presents a plesiomorphic karyotype. This last species showed larger number of chromosomal homeologies with the species of the *Artibeus* genus ($2n = 30/31$), that possess the presumed ancestral karyotype for the subfamily Stenodermatinae (Baker, 1979; Baker *et al.*, 1979; Lim, 1993). The direction of the karyotypic evolution of the genus *Uroderma* trended toward an increase in the chromosome number by events of fission followed by events of fusion, inversion and/or translocation.

Acknowledgments

We would like to thank CAPES, FINEP, CNPq and UFPA for their financial support and the technicians at the Laboratory of Cytogenetics (CCB/UFPA) and MPEG (Museu Paraense Emílio Goeldi) for field work help and technical support.

References

- Baker RJ (1967) Karyotypes of bats of the family Phyllostomidae and their taxonomic implications. *Southwest Nat* 12:407-428.
- Baker RJ (1973) Comparative cytogenetic of the New World leaf-nosed bats (Phyllostomidae). *Period Biol* 75:37-45.
- Baker RJ (1979) Karyology. In: Baker RJ, Jones JK and Carter DC (eds) *Biology of Bats of the New World Family Phyllostomatidae*. Part III. Special Publications of the Museum of Texas Tech University, Texas Tech University Press, Austin, pp 107-155.
- Baker RJ (1981) Chromosomal flow between chromosomally characterized taxa of a volant mammal, *Uroderma bilobatum*. *Evolution* 35:296-305.
- Baker RJ and Lopez G (1970) Chromosomal variation in bats of the genus *Uroderma* (Phyllostomidae). *J Mammal* 51:786-789.
- Baker RJ, Atchley WR and McDaniel VR (1972) Karyology and morphometrics of Peter's tent-making bat, *Uroderma bilobatum* (Chiroptera, Phyllostomatidae). *Syst Zool* 21:414-429.
- Baker RJ, Bleier WJ and Atchley WR (1975) A contact zone between karyotypically characterized taxa of *Uroderma bilobatum* (Mammalia, Chiroptera). *Syst Zool* 24:133-142.
- Baker RJ, Bass RA and Johnson MA (1979) Evolutionary implications of chromosomal homology in four genera of stenodermine bats (Phyllostomidae, Chiroptera). *Evolution* 33:220-226.
- Baker RJ, Haiduk MW, Robbins LW, Cadena A and Koop BF (1982) Chromosomal studies of South American bats and their systematic implications. In: Mares MA and Genoways HH (eds) *Mammalian Biology in South America*. v. 4. Special publication Series, Pymatuning Laboratory of Ecology, University of Pittsburgh, pp 303-327.
- Bredt A, Araújo FAA, Caetano J, Rodrigues MGR, Yoshizawa M, Silva MMS, Massunaga PNT, Bürer SP, Porto VAR and Uieda W (1996) Morcegos em Áreas Urbanas e Rurais: Manual de Manejo e Controle. Fundação Nacional de Saúde, Ministério da Saúde, Brasília, 117 pp.
- Ford CE and Hamerton JL (1956) A colchicine, hypotonic-citrate, squash sequence for mammalian chromosomes. *Stain Technol* 31:247-251.
- Gardner AL (1977) Chromosomal variation in *Vampyressa* and a review of chromosomal evolution in the Phyllostomidae (Chiroptera). *Syst Zool* 26:300-318.
- Hill JE and Smith JD (1984) *Bats. A Natural History*. 2nd edition. University of Texas Press, Austin, Texas, 243 pp.
- Howell WM and Black DA (1980) Controlled silver-staining of nucleolar organizer regions with protective colloidal developer: A 1-step method. *Experientia* 36:1014.
- Lee MR and Elder FFB (1980) Yeast stimulation of bone marrow mitosis for cytogenetic investigations. *Cytogenet Cell Genet* 26:36-40.
- Lim BK (1993) Cladistic reappraisal of neotropical Stenodermatine bat phylogeny. *Cladistics* 9:147-165.
- Nowak RM (1996) *Walker's Bats of the World*. 6^a edition. The Johns Hopkins University Press, Baltimore. 285 pp.
- Owen RD (1987) Phylogenetic analysis of the bat subfamily Stenodermatinae (Mammalia, Chiroptera). Special Publications of the Museum of Texas Tech University, Texas Tech University Press, Lubbock, Texas. 65 pp.
- Rodrigues LRR (1998) *Relações cromossômicas entre Phyllostomus e Artibeus* (Phyllostomidae-Chiroptera). Masters Dissertation, Universidade Federal do Pará, Belém.
- Silva AM (2000) *Estudo citogenético e evolução cromossômica em seis espécies amazônicas da subfamília Stenodermatinae* (Chiroptera, Phyllostomidae). Masters Dissertation, Universidade Federal do Pará, Belém.
- Sumner AT (1972) A simple technique for demonstrating centromeric heterochromatin. *Exp Cell Res* 75:304-306.

- Tucker PK (1986) Sex chromosome-autosome translocations in the leaf-nosed bats, family Phyllostomidae. I. Mitotic analyses of the subfamilies Stenodermatinae and Phyllostominae. *Cytogenet Cell Genet* 43:19-27.
- Tucker PK and Bickham JW (1986) Sex chromosome-autosome translocation in the leaf nosed bats. II. Meiotic analyses of the subfamilies Stenodermatinae and Phyllostominae. *Cytogenet Cell Genet* 43:28-27.
- Van Den Bussche RA (1992) Restriction site variation and molecular systematic of the New World Leaf-nosed bats. *J Mammal* 73:39-42.
- Varella-Garcia M, Morielle-Versute E and Taddei VA (1989) A survey of cytogenetic data on Brazilian bats. *Rev Bras Genet* 12:761-793.

Associate Editor: Yatiyo Yonenaga-Yassuda